


Indicator	Default rate of clients from anti-retroviral therapy, tuberculosis directly observed treatment (TB-DOTS) and prevention of mother-to-child transmission of HIV (PMTCT) programmes	CRF Area	Nutrition	Data Flow	Health Center/Mobile Clinic Patient Register (Client data for ART, TB-DOTs or PMTCT programmes) from WFP
------------------	---	-----------------	-----------	------------------	--



<u>Control</u>	<u>Actions</u>	<u>Responsibility</u>	<u>Frequency</u>
1 Documentation Review	Calculate the % of available, on-time and complete source records and reports that are expected to contain TB-DOTS and HIV-PMTCT programming at the site level, CP Central Office, WFP Field and Country Office.	CO M&E Officer or Assistant or Nutrition Officer	🔄 <i>Frequency Quarterly</i> ★ ★ Highly Recommended
2 Recounting & Reaggregating	N/A	N/A	N/A
3 Recomputing Formulas	Recompute the formula used to use calculate the reported value of the default rate to determine a recomputed value, calculate the % error between the recomputed and reported value, and identify possible reasons for the discrepancy (if any).	CO M&E Officer or Assistant or Nutrition Officer	🔄 <i>Frequency Quarterly</i> ★ Recommended
4 Reconciliation & Triangulation	N/A	N/A	N/A
5 Secondary Sources	N/A	N/A	N/A
6 On-Site Visits	N/A	N/A	N/A
7 Methodological Compliance	N/A	N/A	N/A